


A13

(10)  **Europäisches Patentamt**  
**European Patent Office**  
**Office européen des brevets**

(11) Publication number: **0 208 420**  
**A1**

(12) **EUROPEAN PATENT APPLICATION**

(21) Application number: 86304413.7

(51) Int. Cl.<sup>4</sup>: **C 07 D 417/12**  
**C 07 D 277/34, A 61 K 31/425**

(22) Date of filing: 10.06.86

(30) Priority: 10.06.85 JP 126826/85

(43) Date of publication of application:  
 14.01.87 Bulletin 87/3

(54) Designated Contracting States:  
 AT BE CH DE FR GB IT LI LU NL SE

(71) Applicant: **Takada Chemical Industries, Ltd.**  
 27, Doshomachi 2-chome Higashi-ku  
 Osaka-shi Osaka, 541(JP)


(72) Inventor: **Maguro, Kanji**  
 2-21, Mondosou Nishinomiya  
 Hyogo 662(JP)

(72) Inventor: **Fujita, Takeshi**  
 13-15, Nagasoda 1-chome Takarazuka  
 Hyogo 668(JP)

(74) Representative: **Lewin, John Harvey et al,**  
 Elkington and Fife High Holborn House 52/54 High  
 Holborn  
 London WC1V 6SH(GB)

(54) **Thiazolidine derivatives, their production and use.**

(57) **A thiazolidinedione derivative of the general formula:**



wherein X is an oxygen or sulfur atom, R<sup>1</sup> and R<sup>2</sup> each independently is hydrogen or a hydrocarbon residue which may optionally be substituted and R<sup>1</sup> and R<sup>2</sup> may jointly, together with the oxazole or thiazole ring, form a condensed ring and A is a lower alkylene group; or a salt thereof, are novel compounds, which exhibit in mammals blood sugar- and lipid-lowering activity, and are of value as a therapeutic agent for diabetes and/or therapeutic agent for hyperlipemia.

EP 0 208 420 A1

- 1 -

Thiazolidine Derivatives, Their Production and Use


This invention relates to novel thiazolidinedione derivatives having blood sugar and lipid lowering activity, methods of producing the same and antidiabetic agents containing the same.

Various biguanide type compounds and sulfonylurea type compounds have so far been used as antidiabetic agents. However, biguanides are rather obsolete nowadays because they induce lactic acidosis. Sulfonylureas, on the other hand, have potent blood sugar lowering activity but frequently cause severe hypoglycemia, so that much care is necessary in using them. Accordingly, the appearance of novel type antidiabetics free of these drawbacks has been waited for. Meanwhile, Japanese Patent Publications Kokai No. 22636/80 and Kokai No. 64586/80, Chemical & Pharmaceutical Bulletin, vol. 30, p. 3563 (1982), ibid., vol. 30, p. 3580 (1982) and ibid., vol. 32, p. 2267 (1984) describe that various thiazolidinedione derivatives exhibit blood sugar and lipid lowering activity. Antidiabetic activity of ciglitazone was also reported in Diabetes, 32, P.804 (1983).

The present inventors synthesized and evaluated various compounds not specifically described in the above-cited patent publications and, as a result, compounds having potent pharmacological activity with low toxicity were found.

15           The present invention thus provides:

(1)   Thiazolidinedione derivatives of the general formula:




0208420

wherein X is an oxygen or sulfur atom, R<sup>1</sup> and R<sup>2</sup>  
each independently hydrogen or a hydrocarbon  
residue which may optionally be substituted and  
R<sup>1</sup> and R<sup>2</sup> may jointly, together with the oxazole or  
thiazole ring, form a condensed ring and A is a lower  
alkylene group, and salts thereof;

(2) Pharmaceutical compositions suitable for the  
therapy of a mammal suffering from diabetes and/or  
hyperlipemia, which contain as the effective component  
a thiazolidinedione derivative of the general formula  
(I) or a pharmacologically acceptable salt thereof.

(3) A method of producing thiazolidinedione derivatives  
of the general formula (I) and salts thereof which com-  
prises hydrolyzing a compound of the general formula:




wherein the symbols are as defined above, or a salt

0208420

thereof; and


(4) A method of producing thiazolidinedione derivatives of the general formula:



10 wherein the symbols are as defined above, and salts thereof, which comprises reacting a compound of the general formula:



15 wherein  $R^1$ ,  $R^2$  and X are as defined above and Y is a halogen atom, with a compound of the formula:



25 The hydrocarbon residue represented by  $R^1$  and/or  $R^2$  in the above general formulas (I), (Ia), (II) and (III) may be an aliphatic hydrocarbon residue, an alicyclic hydrocarbon residue, an alicyclic-aliphatic hydrocarbon residue, an aromatic-aliphatic hydrocarbon residue or an aromatic hydrocarbon residue. Said aliphatic hydro-

0208420

carbon residue includes saturated aliphatic hydrocarbon residues containing 1-8 carbon atoms, (e.g. methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, t-butyl, pentyl, hexyl, heptyl and octyl), preferably 1-4 carbon atoms; said alicyclic

5 hydrocarbon residue includes saturated alicyclic hydrocarbon residues containing 3-7, preferably 5-6 carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl, and unsaturated alicyclic hydrocarbon residues containing 5-7 carbon atoms, such as 1-cyclopentenyl,

10 2-cyclopentenyl, 3-cyclopentenyl, 1-cyclohexenyl, 2-cyclohexenyl, 3-cyclohexenyl, 1-cycloheptenyl and 2-cycloheptenyl; said alicyclic-aliphatic hydrocarbon residue includes groups resulting from bonding the above-

15 mentioned alicyclic hydrocarbon residues to the above-mentioned aliphatic hydrocarbon residues and containing 4-9 carbon atoms, such as cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclopentylmethyl, 2-cyclopentenylmethyl, 3-cyclopentenylmethyl, cyclohexylmethyl, 2-cyclohexenylmethyl, cycloheptylmethyl and cycloheptyl-

20 ethyl; said aromatic-aliphatic hydrocarbon residue includes phenylalkyl groups containing 7-9 carbon atoms, such as benzyl, phenethyl, 1-phenylethyl and phenylpropyl, and naphthylalkyl groups containing 11-13 carbon atoms, such as  $\alpha$ -naphthylmethyl,  $\alpha$ -naphthylethyl and  $\beta$ -naphthyl-

25 ethyl; and said aromatic hydrocarbon residue includes, among others, phenyl and naphthyl ( $\alpha$ -naphthyl,  $\beta$ -naphthyl). By saying that  $R^1$  and  $R^2$  jointly, together with the oxazole or thiazole ring, form a condensed ring,

0208420

it is precisely meant that  $R^1$  and  $R^2$ , together with the thiazole or oxazole ring carbon atoms to which they are attaching, form a ring. Thus, it is meant that  $R^1$  and  $R^2$  are combined together to form a saturated or unsaturated divalent chain hydrocarbon residue containing 3-5 carbon atoms.

Examples of said chain hydrocarbon residue are  $-\text{CH}_2\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}=\text{CHCH}_2-$ ,  $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$ ,  $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-\text{CH}_2-$  and  $-\text{CH}=\text{CH}-\text{CH}_2\text{CH}_2\text{CH}_2-$ .

The hydrocarbon residue represented by  $R^1$  and/or  $R^2$  may have at least one substituent in any position thereof. When  $R^1$  and/or  $R^2$  contains an alicyclic group,  $R^1$  and/or  $R^2$  may have, on the ring thereof, 1-3 lower alkyl groups containing 1-3 carbon atoms (e.g. methyl, ethyl, propyl, isopropyl). When  $R^1$  and/or  $R^2$  contains an aromatic hydrocarbon residue or when  $R^1$  and  $R^2$  combinedly form a condensed ring, the ring may be substituted with 1-4 substituents, which may be the same or different. Said substituents include, among others, halogen (fluorine, chlorine, iodine), hydroxy, cyano, trifluoromethyl, lower alkoxy (e.g. one containing 1-4 carbon atoms, such as methoxy, ethoxy, propoxy or butoxy), lower alkyl (e.g. one containing 1-4 carbon atoms, such as methyl, ethyl, propyl, isopropyl or butyl) and lower alkylthio (e.g. one containing 1-3 carbon atoms, such as methylthio, ethylthio, propylthio or isopropylthio).

A lower alkylene group represented by A is a straight or branched chain having 1 to 3 carbon atoms and includes methylene, ethylene, propylene and trimethylene.

The halogen represented by Y in formula (III) includes chlorine, bromine and iodine.

Compounds of the general formula (I) can form salts with bases since they have an acidic nitrogen atom in their thiazolidine ring. Such salts include, among others, pharmacologically acceptable salts such as sodium, potassium, magnesium and calcium salts..

The compounds (I) and salts thereof according to the invention exhibit excellent blood sugar and blood lipid lowering activities in mammals (e.g. mouse, rat, dog, cat, monkey, horse, human) and are low in acute toxicity as well as in subacute toxicity. Therefore, the thiazolidinedione derivatives (I) and salts thereof are useful in the treatment of hyperlipidemia and/or diabetes and complications resulting therefrom. They are generally administered orally in the form of tablets, capsules, powders or granules, for instance. In some instances, they may be administered parenterally in the form of injections, suppositories or pellets, among others. In using them as therapeutic agents for diabetes or hyperlipidemia, they can be administered generally in an oral daily dose of 0.01-10 mg/kg or a parenteral daily dose of 0.005-10 mg/kg. Desirably, they are administered in such dose every day or intermittently 2-4 times a week.

The compounds of general formula (I) or salts thereof can be produced by hydrolyzing the compounds of general formula (II) or salts thereof. This hydrolysis reaction is generally carried out in an appropriate solvent in the presence of water and a mineral acid. Examples of the



solvent which are generally used are alkanols (e.g. methanol, ethanol, propanol, 2-propanol, butanol, isobutanol, 2-methoxyethanol), dimethyl sulfoxide, sulfolane, dioxane and dimethoxyethane. The mineral acid is, for example, hydrochloric acid, hydrobromic acid or sulfuric acid and is used in an amount of 0.1-10 moles, preferably 0.2-3 moles, per mole of the compound of general formula (II). Water is used generally in large molar excess relative to the compound of general formula (II). This reaction is generally carried out with warming or heating and the reaction temperature is generally 60-150°C. The heating time is generally several hours to ten and odd hours.

Those compounds of general formula (I) wherein A is methylene and salts thereof, namely the compounds of general formula (Ia) and salts thereof [hereinafter collectively referred to as "compounds (Ia)"], can be obtained by reacting a compound of general formula (III) with a compound of general formula (IV) or a salt thereof [hereinafter collectively referred to as "compound (IV)"]. The reaction of the compound of general formula (III) and the compound (IV) is generally carried out in the presence of an appropriate solvent and an appropriate base and this reaction gives the compounds (Ia).

Such solvent includes, among others, dimethylformamide, dimethyl sulfoxide, sulfolane, tetrahydrofuran and dimethoxyethane. Examples of said base are sodium hydride, potassium hydride, sodium amide, sodium alkoxide


(e.g. sodium methoxide, sodium ethoxide), potassium alkoxide (e.g. potassium t-butoxide) and potassium carbonate. This reaction is preferably carried out by first allowing formation of a dianion by bringing such  
5 base into contact with the compound (IV) in a molar ratio of 2:1 and thereafter adding the compound of general formula (III) in an amount of 1 mole per mole of compound (IV). This condensation reaction is carried out generally at 0°-120°C, preferably 20°-100°C, and the  
10 reaction time is generally 0.5-5 hours.

The thus-produced thiazolidinedione derivatives (I) and salts thereof can be isolated and purified by known separation/purification techniques such as, for example, concentration, concentration under reduced pressure,  
15 solvent extraction, crystallization, recrystallization, phase transfer and chromatography.

The starting compounds (II) can be produced, for example, by the following methods:

1) The process illustrated below gives those compounds  
20 of general formula (I) wherein X is an oxygen atom, namely the compounds of general formula (IIa), and salts thereof [hereinafter collectively referred to as "compounds (IIa)].

0208420



[In the above formulas, R<sup>1</sup>, R<sup>2</sup> and A are as defined above and R<sup>3</sup> is hydrogen or a lower alkyl group.]

The reaction for deriving the compound (VII) from the compound (V) is carried out in the manner of condensation of the compound (V) with the compound (VI) in the presence of a deacidifying agent (e.g. potassium carbonate, sodium carbonate, sodium hydrogen carbonate, potassium hydrogen carbonate, sodium hydroxide, potas-

0208420

5 sium hydroxide, triethylamine). This reaction can be conducted in a solvent, such as dimethylformamide, dimethyl sulfoxide, tetrahydrofuran, ethyl ether, ethyl acetate, chloroform or dichloromethane, or a mixed solvent prepared by adding water to such solvent as necessary, at  $-10^{\circ}\text{C}$  to  $50^{\circ}\text{C}$ .

10 The compound (VII) is then subjected to ring closure, whereby the compound (VIII) can be derived. This reaction is carried out generally in the presence of a dehydrating agent. Known dehydrating agents, such as phosphorus oxychloride, thionyl chloride, phosphorus pentoxide, polyphosphoric acid, polyphosphoric acid esters, acetic anhydride and sulfuric acid, and mixtures of these, may suitably be used. Although the reaction  
15 conditions may vary depending on the dehydrating agent employed, this reaction can be effected generally in an inert solvent (e.g. benzene, toluene, xylene, dichloromethane, chloroform) at about  $30^{\circ}$ - $140^{\circ}\text{C}$ , or in an excess of the dehydrating agent, which serves also as a solvent, within said temperature range. The dehydrating agent is  
20 used in an amount of 1-30 moles per mole of compound (VII).

25 The reaction for deriving the compound (IX) from the compound (VIII) can be readily carried out in the manner of a conventional catalytic reduction using palladium-on-carbon as catalyst or a conventional reduction using zinc or iron in combination with acetic acid.

0208420

The compound (IX) may be isolated in pure form or may be subjected to the next reaction step without isolation or purification.


5       The reaction for deriving the compound (XI) from the compound (IX) is carried out in the manner of the so-called Meerwein arylation. Thus, the compound (IX) is diazotized in the presence of a hydrohalogenic acid (HY) and then reacted with acrylic acid or an ester thereof (X) in the presence of a copper catalyst (e.g. cuprous  
10       oxide, cupric oxide, cuprous chloride, cupric chloride, cuprous bromide, cupric bromide). The compound (IX) may be purified, for example by chromatography or may be submitted to the next reaction step without isolation and purification.

15       Reaction of the compound (XI) with thiourea then gives (IIa). This reaction is carried out generally in a solvent such as an alcohol (e.g. methanol, ethanol, propanol, 2-propanol, butanol, isobutanol, 2-methoxy-ethanol), dimethylformamide, dimethyl sulfoxide or sulfolane. The reaction temperature is generally 20°-180°C,  
20       preferably 60°-150°C. Thiourea is used in an amount of 1-2 moles per mole of compound (XI). In this reaction, a hydrogen halide is formed as a byproduct with the progress of the reaction, whereby the hydrohalogenic acid salt of compound (IIa) is formed. In this case, the  
25       reaction may be carried out in the presence of sodium acetate, potassium acetate or the like so that the hydro-

0208420

gen halide can be captured thereby and (IIa) can be produced in the free form. Such acid acceptor is used generally in an amount of 1-1.5 moles per mole of compound (XI). Such reaction gives the compound (IIa), which may be isolated as desired or may be submitted directly to the next hydrolysis step without isolation thereof.

2) The compounds (II) can also be produced by the following process:



0208420

by the method described in Chemical & Pharmaceutical Bulletin, vol. 30, p. 3563 (1982).

[Examples]

Example 1

A mixture of 2-imino-5-{4-[2-(4-methyl-5-phenyl-  
5 2-oxazolyl)ethoxy]benzyl}-4-thiazolidinone (5.5 g),  
ethanol (100 ml) and 2 N HCl (60 ml) was heated  
under reflux for 6 hours and then poured into water  
and extracted with chloroform. The chloroform layer  
was washed with water and dried (MgSO<sub>4</sub>). The solvent  
10 was distilled off to give 5-{4-[2-(4-methyl-5-phenyl-  
2-oxazolyl)ethoxy]-2,4-thiazolidinedione (2.8 g,  
50.9%). Recrystallization from ethyl acetate gave  
colorless prisms. M.p. 168-169°C.


Elemental analysis

15 Calcd. for C<sub>22</sub>H<sub>20</sub>N<sub>2</sub>O<sub>4</sub>S: C, 64.69; H, 4.93; N, 6.86  
Found: C, 64.90; H, 5.05; N, 6.82


Examples 2-4

In the same manner as Example 1, the  
compounds listed in Table 1 were obtained.

20 Table 1



0208420

Example No.	R <sup>1</sup>	R <sup>2</sup>	M.P. (°C)	Recrystallization solvent	Yield (%)
2	(CH <sub>3</sub> ) <sub>2</sub> CHCH <sub>2</sub> -	CH <sub>3</sub>	123-124	Ethyl acetate-hexane	88.0
3	H	CH <sub>3</sub>	156-157	Ethanol	71.4
4		CH <sub>3</sub>	175-176	Ethanol-dichloromethane	48.2

#### Example 5

A mixture of 2-imino-5-{4-[2-(5-methyl-4-phenyl-2-oxazolyl)ethoxy]benzyl}-4-thiazolidinone (7.6 g), 1 N H<sub>2</sub>SO<sub>4</sub> (70 ml) and dioxane (70 ml) was stirred at 80°C for 24 hours and then concentrated. The residue was neutralized with potassium carbonate and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried (MgSO<sub>4</sub>). The solvent was then distilled off and the oily residue was subjected to column chromatography using silica gel (120 g). Elution with chloroform-methanol (49:1, v/v) gave 5-{4-[2-(5-methyl-4-phenyl-2-oxazolyl)ethoxy]benzyl}-2,4-thiazolidinedione as an oil (7.06 g, 92.7%).



I R (Neat)  $\text{cm}^{-1}$ : 1755, 1700.

NMR  $\delta$  ppm in  $\text{CDCl}_3$ : 2.48(3H, s), 3.02(1H, d, d, J=14 and 9), 3.21(2H, t, J=7), 3.41(1H, d, d, J=14 and 4), 4.35(2H, t, J=7), 4.41(1H, d, d, J=9 and 4), 5 6.84(2H, d, J=9), 7.11(2H, d, J=9), 7.2~7.75(5H, m), 9.30(1H, broad).

The above-obtained oily substance (7.0 g) was dissolved in methanol (50 ml), and 5 N NaOMe (MeOH solution, 3.77 ml) was added to the solution. The mixture was stirred at room temperature for 10 minutes and concentrated and, then, the residue was treated with ether to give sodium salt of 5-[4-[2-(5-methyl-4-phenyl-2-oxazolyl)ethoxy]benzyl]-2,4-thiazolidine-15 dione (5.8 g, 78.6%). Recrystallization from methanol-ether gave colorless prisms. M.p. 261-262°C (decomposition).

#### Elemental analysis

Calcd. for  $\text{C}_{22}\text{H}_{19}\text{N}_2\text{O}_4\text{SNa}$ : C, 61.39; H, 4.45; N, 6.51  
20 Found : C, 61.56; H, 4.56; N, 6.64

#### Example 6

The procedure of Example 5 was followed to give sodium salt of 5-[4-[2-(5-ethyl-4-phenyl-2-oxazolyl)-ethoxy]benzyl]-2,4-thiazolidinedione in 76.9% yield. 25 Recrystallization from ethanol gave colorless prisms. M.p. 248-250°C (decomposition).

Elemental analysis

Calcd. for  $C_{23}H_{21}N_2O_4SNa$ : C, 62.15; H, 4.76; N, 6.30

Found : C, 61.76; H, 4.66; N, 6.40

Example 7

- 5 Sodium hydride (60% in oil, 1.2 g) was added to a solution of 5-(4-hydroxybenzyl)-2,4-thiazolidinedione (3.4 g) in DMF (30 ml), and the mixture was stirred at room temperature for 30 minutes, followed by dropwise addition of a solution of 2-chloromethyl-4-phenylthiazole
- 10 (4.4 g) in DMF (20 ml) at room temperature. The mixture was stirred at room temperature for 1 hour and at 60°C for 1 hour, poured into water, neutralized with acetic acid and extracted with ethyl acetate. The ethyl acetate layer was washed with water and
- 15 dried ( $MgSO_4$ ). The solvent was then distilled off and the oily residue was subjected to column chromatography using silica gel (100 g). Elution with benzene-acetone (25:1, v/v) gave 5-[4-(4-phenyl-2-thiazolylmethoxy)benzyl]-2,4-thiazolidine-
- 20 dione (2.9 g, 49.2%). Recrystallization from ethanol gave light-yellow crystals. M.p. 164-165°C.

Elemental analysis


Calcd. for  $C_{20}H_{16}N_2O_3S_2$ : C, 60.59; H, 4.07; N, 7.07

Found : C, 60.67; H, 4.03; N, 7.14

Examples 8-9

The procedure of Example 7 was repeated to give the compounds listed in Table 2.

Table 2



Example No.	X	M.P. (°C)	Recrystallization solvent	Yield (%)
8	O	188-189	Dichloromethane-methanol	32.5
9	S	184-185	Dichloromethane-methanol	35.3

Example 10

Production of tablets

- a) (1) Sodium salt of 5-[4-[2-(5-methyl-4-phenyl-2-oxazolyl)ethoxy]benzyl]-2,4-thiazolidinedione 30 g
- (2) Lactose 50 g
- (3) Corn starch 15 g
- (4) Carboxymethylcellulose calcium 44 g
- (5) Magnesium stearate 1 g
- 1,000 tablets 140 g

A mixture of the indicated quantities of (1), (2) and (3), 30 g of (4) and an adequate quantity of water

is kneaded, then dried under vacuum, and granulated. The granular composition obtained is mixed with 14 g of (4) and 1 g of (5) and the resulting mixture is tableted on a tableting machine to give 1,000 tablets each containing 30 mg of (1).

Reference Example 1

A mixture of 3-(4-nitrophenoxy)propionic acid (10.5 g), thionyl chloride (11.9 g), N,N-dimethylformamide (0.3 g) and toluene (100 ml) was stirred at 90°C for 1 hour, then concentrated under reduced pressure, and the oily residue was dissolved in ethyl acetate (30 ml). The solution was added dropwise to a mixture of 3-amino-5-methyl-2-hexanone hydrochloride (8.3 g), sodium carbonate (10.6 g), water (200 ml) and ethyl acetate (100 ml) at room temperature. The mixture was stirred at room temperature for 1 hour and the ethyl acetate layer was separated. The ethyl acetate layer was washed with water and dried (MgSO<sub>4</sub>). The solvent was distilled off to give 3-[3-(4-nitrophenoxy)-propionylamino]-5-methyl-2-hexanone (11.5 g, 71.4%). Recrystallization from ethyl acetate-hexane gave colorless prisms. M.p. 101-102°C.


Elemental analysis





Calcd. for C<sub>16</sub>H<sub>22</sub>N<sub>2</sub>O<sub>5</sub>: C, 59.62; H, 6.88; N, 8.69

Found : C, 59.67; H, 6.79; N, 8.61

In the same manner as above, there were obtained the compounds listed in Table 3.

Table 3



R <sup>1</sup>	R <sup>2</sup>	M.P. (°C)	Recrystallization solvent	Yield (%)
CH <sub>3</sub>		134-135	Ethyl acetate-hexane	75.5
H	CH <sub>3</sub>	147-148	Ethanol	52.6
	CH <sub>3</sub>	131-132	Ethyl acetate	79.6
	CH <sub>3</sub>	143-144	Ethyl acetate-hexane	63.2
	C <sub>2</sub> H <sub>5</sub>	132-133	Ethyl acetate	76.2

Reference Example 2

A mixture of 3-[3-(4-nitrophenoxy)propionylamino]-5-methyl-2-hexanone (11.0 g), phosphorus oxychloride (6.3 g) and toluene (100 ml) was stirred under reflux for 1 hour. After cooling, the mixture was poured into ethyl acetate (200 ml), washed with saturated aqueous sodium hydrogen carbonate and water in that order and dried (MgSO<sub>4</sub>). The solvent was then distilled off and the oily residue was subjected to column chromatography using silica gel (150 g), whereby

4-isobutyl-5-methyl-2-[2-(4-nitrophenoxy)ethyl]oxazole  
(8.1 g, 78.6%) was recovered from an ethyl acetate-  
hexane (1:4, v/v) eluate fraction. Recrystallization from  
ether-hexane gave colorless needles. M.p. 45-46°C.

5 Elemental analysis


Calcd. for  $C_{16}H_{20}N_2O_4$ : C, 63.14; H, 6.62; N, 9.20

Found : C, 63.14; H, 6.44; N, 9.27

In the same manner as above, there were obtained  
the following compounds listed in Table 4.

10

Table 4



15

R <sup>1</sup>	R <sup>2</sup>	M. P. (°C)	Recrystallization solvent	Yield (%)
CH <sub>3</sub>		117-118	Ethyl acetate-hexane	89.0
H	CH <sub>3</sub>	89-90	Ethyl acetate-hexane	82.3
	C <sub>2</sub> H <sub>5</sub>	72-73	Ether-hexane	89.5
	CH <sub>3</sub>	91-92	Ether-hexane	81.7
	CH <sub>3</sub>	88-89	Ether-hexane	86.3

20

Reference Example 3

A solution of 4-methyl-2-[2-(4-nitrophenoxy)ethyl]-  
5-phenyloxazole (12.5 g) in methanol (150 ml) was

25

subjected to catalytic reduction in the presence of  
5% Pd-C (wet, 3.0 g). The catalyst was then filtered  
off and the filtrate was concentrated to give 2-[2-(4-  
aminophenoxy)ethyl]-4-methyl-5-phenyloxazole (11.0 g,  
5 97.3%). Recrystallization from ethanol gave colorless  
needles. M.p. 106-107°C.

Elemental analysis

Calcd. for  $C_{18}H_{18}N_2O_2$ : C, 73.45; H, 6.16; N, 9.52

Found : C, 73.53; H, 6.11; N, 9.44

10 The above procedure was followed to give 2-[2-  
(4-aminophenoxy)ethyl]-5-ethyl-4-phenyloxazole in  
98.7% yield. Recrystallization from ether-hexane  
gave colorless prisms. M.p. 89-90°C.

Elemental analysis

15 Calcd. for  $C_{19}H_{20}N_2O_2$ : C, 74.00; H, 6.54; N, 9.08

Found : C, 74.05; H, 6.28; N, 9.25

Reference Example 4

1) 2-[2-(4-Aminophenoxy)ethyl]-4-methyl-5-phenyl-  
oxazole (10.5 g) was dissolved in acetone (100 ml)-  
20 methanol (30 ml), and 47% aqueous HBr (24.6 g) was  
added to the solution, followed by dropwise addition  
of a solution of  $NaNO_2$  (2.7 g) in water (10 ml) at 5°C.  
The mixture was stirred at 5°C for 15 minutes and methyl  
acrylate (18.4 g) was added thereto. The resulting  
25 mixture was then warmed to 38°C and, with stirring

vigorously, cuprous oxide powder (1 g) was added portionwise to the mixture. The mixture was stirred until completion of nitrogen gas generation, then concentrated, and the residue was made basic with aqueous ammonia and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried ( $\text{MgSO}_4$ ). The solvent was distilled off to give crude 2-bromo-3-{4-[2-(4-methyl-5-phenyl-2-oxazolyl)ethoxy]-phenyl}propionate as an oil (14.3 g, 89.9%).

I R (Neat)  $\text{cm}^{-1}$ : 1740.

NMR  $\delta$  ppm in  $\text{CDCl}_3$ : 2.37(3H.s). 3.0~3.6(2H.m). 3.25(2H.t,  $J=7$ ). 3.67(3H.s). 4.2~4.5(3H.m). 6.7~7.7(9H.m).

2) To a solution of the oily substance (14.0 g) obtained in the above procedure 1) in ethanol (150 ml) were added thiourea (2.4 g) and sodium acetate (2.6 g), and the mixture was stirred under reflux for 3 hours and then concentrated. The residue was neutralized with saturated aqueous sodium hydrogen carbonate, followed by addition of ether (50 ml)-hexane (50 ml). The mixture was stirred for 10 minutes and the resulting crystalline precipitate was collected by filtration to give 2-imino-



0208420

5-{4-[2-(4-methyl-5-phenyl-2-oxazolyl)ethoxy]benzyl}-  
4-thiazolidinone (6.0 g, 46.2%). Recrystallization  
from chloroform-methanol gave colorless prisms. M.p.  
194-195°C.

5 Elemental analysis

Calcd. for  $C_{22}H_{21}N_3O_3S$ : C, 64.85; H, 5.19; N, 10.31  
Found : C, 64.67; H, 5.03; N, 10.02

In the same manner as above, there was obtained  
5-{4-[2-(5-ethyl-4-phenyl-2-oxazolyl)ethoxy]benzyl}-  
10 2-imino-4-thiazolidinone. The overall yield from  
the corresponding amino compound was 39.2%. Recrystal-  
lization from methanol gave colorless prisms. M.p.  
164-165°C.

Elemental analysis

15 Calcd. for  $C_{23}H_{23}N_3O_3S$ : C, 65.54; H, 5.50; N, 9.97  
Found : C, 65.32; H, 5.42; N, 9.95

Reference Example 5

A solution of 4-isobutyl-5-methyl-2-[2-(4-nitro-  
phenoxy)ethyl]oxazole (7.8 g) in methanol (100 ml) was  
20 subjected to catalytic reduction in the presence of  
5% Pd-C (wet, 2.0 g). The catalyst was then filtered  
off and the filtrate was concentrated to give an amino  
compound as an oil. The amino compound was dissolved  
in acetone (50 ml)-methanol (20 ml), and 47% aqueous  
25 HBr (17.9 g) was added, followed by dropwise addition

0208420

of a solution of  $\text{NaNO}_2$  (1.9 g) in water (6 ml) at  $5^\circ\text{C}$  or lower. The mixture was stirred at  $5^\circ\text{C}$  for 15 minutes. Methyl acrylate (15.7 g) was then added and the resulting mixture was warmed to  $38^\circ\text{C}$  and, with stirring  
5 vigorously, cuprous oxide powder (0.5 g) was added portionwise to the mixture. The mixture was stirred until completion of nitrogen gas generation and then concentrated. The residue was made basic with aqueous ammonia and extracted with ethyl acetate. The ethyl  
10 acetate layer was washed with water and dried ( $\text{MgSO}_4$ ). The solvent was distilled off to give crude methyl 2-bromo-3-[4-[2-(4-isobutyl-5-methyl-2-oxazolyl)ethoxy]phenyl]-propione as an oil (9.5 g, 86.4%).

I R (Neat)  $\text{cm}^{-1}$ : 1740.

15 NMR  $\delta$  ppm in  $\text{CDCl}_3$ : 0.88(6H.d.J=7), 1.8-2.1(1H.m), 2.17(3H.s), 2.2-2.4(2H.m), 3.0-3.5(2H.m), 3.12(2H.t.J=7), 3.68(3H.s), 4.2-4.5(1H.m), 4.28(2H.t.J=7), 6.7-7.4(4H.m)

20 2) To a solution of the oily product (9.2 g) obtained in the above procedure 1) in ethanol (100 ml) were added thiourea (1.7 g) and sodium acetate (1.8 g), and the mixture was stirred under reflux for 3 hours and then concentrated. The residue was neutralized with  
25 saturated aqueous sodium hydrogen carbonate, followed

0208420

by addition of ether (50 ml)-hexane (50 ml). The mixture was stirred for 10 minutes and the resulting crystalline precipitate was collected by filtration and recrystallized from ethyl acetate to give 2-imino-5-{4-[2-(4-isobutyl-5-methyl-2-oxazolyl)ethoxy]benzyl}-4-thiazolidinone (3.0 g, 35.7%) as colorless prisms. M.p. 167-168°C.


Elemental analysis



Calcd. for  $C_{20}H_{25}N_3O_3S$ : C, 61.99; H, 6.50; N, 10.84

Found : C, 61.96; H, 6.39; N, 10.70

In the same manner as above, there were obtained the compounds listed in Table 6. The yield is the overall yield from the corresponding starting amino compound.

Table 6



R <sup>1</sup>	R <sup>2</sup>	M.P. (°C)	Recrystallization solvent	Yield (%)
H	CH <sub>3</sub>	198-199	Methanol	38.6
	CH <sub>3</sub>	152-154	Ethyl acetate	50.1
	CH <sub>3</sub>	178-180	Methanol	47.2

0208420

Test Example

Blood sugar and lipid lowering activities in mice

The test compound was mixed with a powder diet (CE-2, Clea Japan) at an addition level of 0.005% and the diet was given to KKA<sup>y</sup> mice (male, 8-10 weeks old; 5 mice per group) ad libitum for 4 days, during which the mice were freely accessible to water. Blood samples were collected from the orbital venous plexus and assayed for blood sugar level by the glucose oxidase method and for plasma triglyceride (TG) level by enzymatically determining glycerol formed using Cleantech TG-S kit (Iatron). Both the activity levels were calculated using the formula given below. The results thus obtained are shown in Table 7. For comparison, data for a known compound of a similar structure are also given.

$$\frac{(\text{Level in untreated group}) - (\text{Level in treated group})}{(\text{Level in untreated group})} \times 100$$

0208420

Table 7

	Blood sugar (%) lowering activity	TG lowering activity (%)
1	18**	7
2	53****	43****
4	54****	51****
5	54****	60****
6	46****	51****
7	43****	42****
8	35**	24
9	43****	43****
control compound 1) ciglitazone	10	-13

t-Test \*\*p<0.05. \*\*\*p<0.02. \*\*\*\*p<0.01.

\*\*\*\*\*p<0.001


1) 5-[4-(1-Methylcyclohexylmethoxy)]

benzyl-2,4-thiazolidinedione

0208420

1 Claims

1. A thiazolidinedione derivative of the general formula:



wherein X is an oxygen or sulfur atom, R<sup>1</sup> and R<sup>2</sup> each independently is hydrogen or a hydrocarbon residue which may optionally be substituted and R<sup>1</sup> and R<sup>2</sup> may jointly, together with the oxazole or thiazole ring, form a condensed ring and A is a lower alkylene group; or a salt thereof.

2. A compound as claimed in claim 1, wherein R<sup>1</sup> is a saturated aliphatic hydrocarbon residue containing 1 to 8 carbon atoms, a saturated alicyclic hydrocarbon residue containing 3 to 7 carbon atoms, a phenyl or a naphthyl.

3. A compound as claimed in claim 1, wherein R<sup>2</sup> is a saturated aliphatic hydrocarbon residue containing 1 to 8 carbon atoms, a phenyl or a naphthyl.

4. A compound as claimed in claim 1, wherein R<sup>1</sup> and R<sup>2</sup> jointly, together with the oxazole or thiazole ring, form a condensed ring.

5. A compound as claimed in claim 1, wherein X is an oxygen.

6. A compound as claimed in claim 1, wherein X is sulfur.

7. A compound as claimed in claim 1, wherein A is methylene.


0208420

8. A compound as claimed in claim 1, wherein the compound is 5-{4-[2-(4-isobutyl-5-methyl-2-oxazolyl)ethoxy]benzyl}-2,4-thiazolidinedione.

9. A compound as claimed in claim 1, wherein the compound is 5-{4-[2-(4-cyclohexyl-5-methyl-2-oxazolyl)ethoxy]benzyl}-2,4-thiazolidinedione.


10. A compound as claimed in claim 1, wherein the compound is 5-{4-[2-(5-methyl-4-phenyl-2-oxazolyl)ethoxy]benzyl}-2,4-thiazolidinedione.

11. A pharmaceutical composition suitable for the therapy of a mammal suffering from diabetes and/or hyperlipemia which contains as the effective component a thiazolidinedione derivative of the general formula:




wherein X is an oxygen or sulfur atom, R<sup>1</sup> and R<sup>2</sup> each independently is hydrogen or a hydrocarbon residue which may optionally be substituted and R<sup>1</sup> and R<sup>2</sup> may jointly, together with the oxazole or thiazole ring, form a condensed ring and A is a lower alkylene group, or a pharmacologically acceptable salt thereof.

12. A method of producing a thiazolidinedione derivative of the general formula:




0208420

wherein X is an oxygen or sulfur atom, R<sup>1</sup> and R<sup>2</sup> each independently is hydrogen or a hydrocarbon residue which may optionally be substituted and R<sup>1</sup> and R<sup>2</sup> may jointly, together with the oxazole or thiazole ring, form a condensed ring and A is a lower alkylene group, or a salt thereof, which comprises hydrolyzing a compound of the general formula:




wherein the symbols are as defined above, or a salt thereof.

13. A method of producing a thiazolidinedione derivative of the general formula:




wherein X is an oxygen or sulfur atom, R<sup>1</sup> and R<sup>2</sup> each independently is hydrogen or a hydrocarbon residue which may optionally be substituted and R<sup>1</sup> and R<sup>2</sup> may jointly, together with the oxazole or thiazole ring, form a condensed ring, or a salt thereof, which comprises reacting a compound of the general formula:





0208420


wherein Y is a halogen atom and the other symbols are as defined above, with a compound of the formula:




0208420

Claims (for the Contracting State AT):

1. A method of producing a thiazolidinedione derivative of the general formula:




wherein X is an oxygen or sulfur atom, R<sup>1</sup> and R<sup>2</sup> each independently is hydrogen or a hydrocarbon residue which may optionally be substituted and R<sup>1</sup> and R<sup>2</sup> may jointly, together with the oxazole or thiazole ring, form a condensed ring and A is a lower alkylene group, or a salt thereof, which comprises hydrolyzing a compound of the general formula:



wherein the symbols are as defined above, or a salt thereof.

2. A method of producing a thiazolidinedione derivative of the general formula:




0208420

- 1 wherein X is an oxygen or sulfur atom, R<sup>1</sup> and R<sup>2</sup> each  
independently is hydrogen or a hydrocarbon residue  
which may optionally be substituted and R<sup>1</sup> and R<sup>2</sup> may  
jointly, together with the oxazole or thiazole ring,  
5 form a condensed ring, or a salt thereof, which comprises  
reacting a compound of the general formula:




- 15 wherein Y is a halogen atom and the other symbols are  
as defined above, with a compound of the formula:



- 25 3. A method as claimed in claim 1 or 2, wherein R<sup>1</sup> is  
a saturated aliphatic hydrocarbon residue containing  
1 to 8 carbon atoms, a saturated alicyclic hydrocarbon  
residue containing 3 to 7 carbon atoms, a phenyl or  
a naphthyl.
- 30 4. A method as claimed in any of claims 1 to  
3, wherein R<sup>2</sup> is a saturated aliphatic hydrocarbon  
residue containing 1 to 8 carbon atoms, a phenyl or  
a naphthyl.
- 35 5. A method as claimed in claim 1 or 2, wherein  
R<sup>1</sup> and R<sup>2</sup> jointly, together with the oxazole or thiazole  
ring, form a condensed ring.

- 1 6. A method as claimed in any of claims 1 to 5, where in X is an oxygen.
7. A method as claimed in any of claims 1 to 5, wherein X is sulfur.
8. A method as claimed in any of claims 1 and 3 to 7, wherein A is methylene.
- 10 9. A method as claimed in claim 1 or 2, wherein the derivative is 5-{4-[2-(4-isobutyl-5-methyl-2-oxazolyl)-ethoxy]benzyl}-2,4-thiazolidinedione.
- 15 10. A method as claimed in claim 1 or 2, wherein the derivative is 5-{4-[2-(4-cyclohexyl-5-methyl-2-oxazolyl)ethoxy]benzyl}-2,4-thiazolidinedione.
11. A method as claimed in claim 1 or 2, wherein the derivative is 5-{4-[2-(5-methyl-4-phenyl-2-oxazolyl)-ethoxy]benzyl}-2,4-thiazolidinedione.
- 20 12. A pharmaceutical composition suitable for the therapy of a mammal suffering from diabetes and/or hyperlipemia which contains as the effective component a thiazolidinedione derivative of the general formula:
- 25



wherein X is an oxygen or sulfur atom,  $\text{R}^1$  and  $\text{R}^2$  each independently is hydrogen or a hydrocarbon residue which may optionally be substituted and  $\text{R}^1$  and  $\text{R}^2$  may

jointly, together with the oxazole or thiazole ring, form a condensed ring and A is a lower alkylene group, or a pharmacologically acceptable salt thereof.

5

10

25

30

35



European Patent  
Office

# EUROPEAN SEARCH REPORT

**0208420**  
Application number

EP 86 30 4413

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
A	EP-A-O 139 421 (SANKYO)		C 07 D 417/12 C 07 D 277/34 A 61 K 31/425
P, A	--- EP-A-O 177 353 (TAKEDA) -----		
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
			C 07 D 417/00 C 07 D 277/00 A 61 K 31/00
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 15-10-1986	Examiner DE BUYSER I.A.F.
<b>CATEGORY OF CITED DOCUMENTS</b>			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : member of the same patent family, corresponding document	